

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 31-10-2005		2. REPORT TYPE Journal Article		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Theory of Monte-Carlo Simulations of the Magnetic Circular Dichroism (MCD) Spectra of Alkali Metal/Rare Gas Systems (POSTPRINT)				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) John W. Kenney (Concordia University); Jerry A. Boatz (AFRL/PRSP); Heidi A. Terrill Vosbein (Stennis Space Center)				5d. PROJECT NUMBER 23030423	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRSP 10 E. Saturn Blvd. Edwards AFB CA 93524-7680				8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-PR-ED-JA-2005-439	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S) AFRL-PR-ED-JA-2005-439	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited (AFRL-ERS-PAS-2005-277)					
13. SUPPLEMENTARY NOTES ©2005 Wiley Periodicals, Inc., Published in the International Journal of Quantum Chemistry, Vol. 103, 854-865 (2005).					
14. ABSTRACT The history of magnetic circular dichroism (MCD) spectroscopy in the study of alkali metal/rare gas (M/Rg) cryogenic systems is reviewed in the context of developing a better understanding of alkali metal/hydrogen systems of current interest to the U.S. Air Force as enhanced-performance cryogenic rocket propellants. A new theory for simulating the MCD spectra of M/Rg systems is presented together with a careful discussion of the theory's implicit and explicit approximations and their implications.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT A	18. NUMBER OF PAGES 13	19a. NAME OF RESPONSIBLE PERSON Dr. Scott A. Shackelford
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (include area code) N/A

Theory of Monte Carlo Simulations of the Magnetic Circular Dichroism Spectra of Alkali Metal/Rare Gas Systems

JOHN W. KENNEY, III,¹ JERRY A. BOATZ,²
HEIDI A. TERRILL VOSBEIN³

¹Chemical Physics Laboratory, Concordia University, 1530 Concordia West, Irvine, California 92612

²Air Force Research Laboratory, Space and Missile Propulsion Division; AFRL/PRSP, 10 E. Saturn Blvd., Edwards AFB, California 93524

³Naval Research Laboratory, Acoustic Simulation, Measurements and Tactics Branch, Code 7180, Bldg. 1005, Stennis Space Center, Mississippi 39529

Received 3 February 2004; accepted 12 May 2004

Published online 9 March 2005 in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/qua.20270

ABSTRACT: The history of magnetic circular dichroism (MCD) spectroscopy in the study of alkali metal/rare gas (M/Rg) cryogenic systems is reviewed in the context of developing a better understanding of alkali metal/hydrogen systems of current interest to the U. S. Air Force as enhanced-performance cryogenic rocket propellants. A new theory for simulating the MCD spectra of M/Rg systems is presented together with a careful discussion of the theory's implicit and explicit approximations and their implications. This theory uses a classical Monte Carlo (MC) simulation scheme to model the perturbing effects of the Rg environment on the $^2S \rightarrow ^2P$ MCD-active transition of the M atom. The theory sets up the MC-MCD simulation as a 6×6 matrix eigenvalue/eigenvector problem in the 2P manifold in which are included the effects of M-Rg interactions, metal atom spin-orbit coupling in the 2P manifold, magnetic Zeeman perturbations of the 2S and 2P manifolds, Boltzmann temperature factors, and electric dipole transition moment integrals for left circularly polarized (LCP) and right circularly polarized (RCP) light. The theory may be applied to any type of trapping site of the host M in the guest Rg matrix; a single atom substitutional metal atom trapping

Correspondence to: H. A. Terrill Vosbein; e-mail: terrill@nrlssc.navy.mil

Contract grant sponsors: Air Force Office of Scientific Research-Summer Faculty Research Program (AFOSR-SFRP); Graduate Student Research Program (AFOSR-GSRP); Research Corporation Partners in Science Program.

site (one host Rg atom is replaced by one guest M atom) is modeled in this study for $M = \text{Na}$ and $\text{Rg} = \text{Ar}$. Two temperature factors are used in these simulations; a lattice temperature to model the mobility of the Rg lattice and a magnetic temperature to model Boltzmann factors in the 2S ground manifold. The 6×6 eigenvalue/eigenvector problem is solved for a number of randomly generated and suitably averaged Rg configurations to yield the simulated MC-MCD spectrum for the single substitutional Na/Ar system. The MC-MCD simulations of Na/Ar give the characteristic triplet MCD spectrum with the correct Boltzmann temperature dependence. The simulated MC-MCD spectrum correctly inverts when the direction of the applied magnetic field is reversed. Addition of the LCP and RCP absorbances gives rise to a characteristic $^2S \rightarrow ^2P$ triplet absorption feature. © 2005 Wiley Periodicals, Inc. *Int J Quantum Chem* 103: 854–865, 2005

Introduction

Alkali metal/rare gas solids (M/Rg solids) constitute an important class of model systems of interest to the U. S. Air Force in simulating the properties of enhanced-performance cryogenic rocket propellants [1–4]. In particular, the alkali metal/rare gas (M/Rg) systems share many physical, chemical, and spectroscopic characteristics with M/hydrogen systems, where in a potential rocket propulsion application, M is a high-energy density material (HEDM), e.g., a low mass alkali metal or Group 13 or 14 element [5, 6]. Boatz and Fajardo [7] recently modeled $^2S \rightarrow ^2P$ electronic absorption spectra in M/Rg systems by using quantum mechanical first-order degenerate perturbation theory in conjunction with a classical Monte Carlo (MC) method to account statistically for the effects of the Rg perturbations on the 2S and 2P terms of the M/Rg systems. Their simulations, while not completely quantitative, do recover many of the key attributes of the electronic absorption spectra of these systems, including the characteristic three-peaked or “triplet” absorption band shape. Lawrence and Apkarian [8] take a very similar approach in modeling the 2P terms of halogen atoms doped in cryogenic matrices (X/Rg). Boatz and Fajardo make the reasonable assumption that spin-orbit coupling represents a small perturbation of the 2P term compared with the Rg perturbation, at least for the lighter alkali metals of interest to them as HEDM rocket propellant dopants, and on this basis justify the exclusion of 2P spin-orbit coupling terms in their model. The Boatz and Fajardo first-order perturbation matrix for the 2P term is, therefore a 3×3 matrix whose matrix elements are defined with respect to a suitable zeroth-order orbital p basis in

the angular momentum representation ($|n l m_l\rangle = |n 1 1\rangle, |n 1 0\rangle, |n 1 -1\rangle$). Spin is not considered. Lawrence and Apkarian do include a 2P spin-orbit term for the halogen in their treatment. They argue convincingly that the large halogen 2P spin-orbit coupling interaction is not greatly affected by the matrix environment; i.e., the use of the atomic spin-orbit coupling constant of the halogen in the formalism represents a good approximation. The Lawrence and Apkarian 2P perturbation matrix is a 6×6 matrix whose individual matrix elements are defined with respect to a suitable zeroth-order Cartesian p basis with spin $\{|n l e s m_s\rangle = |n l x \frac{1}{2} \frac{1}{2}\rangle, |n l y \frac{1}{2} \frac{1}{2}\rangle, |n l z \frac{1}{2} \frac{1}{2}\rangle, |n l x \frac{1}{2} -\frac{1}{2}\rangle, |n l y \frac{1}{2} -\frac{1}{2}\rangle, |n l z \frac{1}{2} -\frac{1}{2}\rangle\}$.

Magnetic circular dichroism (MCD) spectroscopy, which measures the differential absorption of left circularly polarized (LCP) light vs. right circularly polarized (RCP) light in a sample placed in a magnetic field aligned parallel to the propagation direction of the light, has a long and venerable history of providing insights to the nature of the chemical and physical environment surrounding the MCD-active chromophore [9]. In particular, the MCD-active $^2S \rightarrow ^2P$ electronic transitions of alkali halide F centers [10–14] and M/Rg systems [15–18] have attracted considerable experimental and theoretical attention over the years. The usual approach is to extract parameters such as the g -value, the spin-orbit coupling constant, and linear vibrational coupling constants from a moment analysis of the experimental MCD spectrum (or spectra) [9, 11]. Only a few full MCD spectral simulations exist in the literature. These simulations of $^2S \rightarrow ^2P$ MCD spectra rely on dynamic Jahn–Teller models in which an a priori choice is made as to which specific interaction vibrational modes will act to lift the degeneracy of the 2P term [19, 20]. This simulation approach is of limited value in modeling the MCD

spectra of M atoms trapped in novel and heretofore unobserved sites in Rg matrices for which specific site geometries and lattice mode types are as yet unknown. A MC approach for handling Rg perturbations in MCD simulations of M/Rg systems naturally suggests itself. A full development of the theory of Monte Carlo-Magnetic Circular Dichroism (MC-MCD) simulations of $^2S \rightarrow ^2P$ MCD spectra is presented in the sections that follow. The MC-MCD simulation method is a first-order degenerate perturbation method that straightforwardly extends the MC absorption-simulation methods of Boatz and Fajardo [7] and Lawrence and Apkarian [8] by (i) including both spin-orbit and Zeeman perturbations in the Hamiltonian, and (ii) separately treating the $^2S \rightarrow ^2P$ transitions associated with LCP and RCP light. The MC-MCD perturbation matrix for the 2P term is a 6×6 matrix whose individual matrix elements are expressed in terms of a suitable uncoupled angular momentum p basis set with spin

$$\{|n l m_l s m_s\rangle = |n 1 1 \frac{1}{2} \frac{1}{2}\rangle, |n 1 0 \frac{1}{2} \frac{1}{2}\rangle, |n 1 -1 \frac{1}{2} \frac{1}{2}\rangle, \\ |n 1 1 \frac{1}{2} -\frac{1}{2}\rangle, |n 1 0 \frac{1}{2} -\frac{1}{2}\rangle, |n 1 -1 \frac{1}{2} -\frac{1}{2}\rangle\}.$$

Theory

As noted above, the theory of MC-MCD spectroscopy of $^2S \rightarrow ^2P$ excitations in M/Rg systems represents an extension of the formalism for MC-absorption spectroscopy developed by Boatz and Fajardo [7] (for M/Rg systems) and by Lawrence and Apkarian [8] (for X/Rg systems). Key aspects of both the Boatz and Fajardo and Lawrence and Apkarian notational schemes are adopted in the following formal development.

TOTAL HAMILTONIAN AND ZERO-ORDER EIGENVECTORS/EIGENVALUES

The total Hamiltonian for an alkali metal M atom imbedded in a cryogenic rare gas matrix composed of N Rg atoms in the presence of an applied external magnetic field is given by

$$H = H_M(\mathbf{r}) + H_{\text{int}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N), \quad (1)$$

where

$$H_M(\mathbf{r}) = T_M(\mathbf{r}) + V_M(\mathbf{r}) \quad (2)$$

is the one-electron Hamiltonian for the optically active electron on M at position \mathbf{r} , composed of a kinetic energy component $T_M(\mathbf{r})$ and an electrostatic potential energy component $V_M(\mathbf{r})$. The potential $V_M(\mathbf{r})$ is the potential experienced by the optically active electron in the free gas-phase M atom in the absence of spin-orbit coupling or an external magnetic field.

To describe completely the 2S ground term of M, two zeroth-order eigenvectors $\{|n 0 0 \frac{1}{2} m_s\rangle; m_s = 1/2, -1/2\}$ are needed; for the 2P excited term, six zeroth-order eigenvectors $\{|n l m_l s m_s\rangle; m_l = 1, 0, -1, m_s = 1/2, -1/2\}$ are required. An uncoupled angular momentum representation $\{|n l m_l s m_s\rangle\}$ for the zeroth-order eigenvectors is chosen, without loss of generality, because of the transparent way in which MCD selection rules may be expressed in terms of this representation (vide infra). However, it is equally valid to use zeroth-order eigenvector sets expressed in the coupled angular momentum representation $\{|n l s j m\rangle\}$ or the Cartesian representation $\{|n l e s m_s\rangle\}$ ($e = x, y, z$ for $l = 1$), since the uncoupled, coupled, and Cartesian representations are mutually related by unitary transformations [21]. In this treatment, the interaction Hamiltonian is composed of four terms, each of which depends on the nuclear position vectors $\{\mathbf{R}_k\}$ of all of the N Rg atoms in the matrix; a metal-rare gas interaction term, a rare gas-rare gas interaction term, a spin-orbit term, and a Zeeman term,

$$H_{\text{int}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = V_{\text{M-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) \\ + V_{\text{Rg-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) + H_{\text{SO}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) \\ + H_{\text{Zeeman}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N). \quad (3)$$

The M nucleus is taken to be the origin of the coordinate system with respect to which \mathbf{r} and the $\{\mathbf{R}_k\}$ are defined. In this treatment, it is assumed, following Lawrence and Apkarian, that M-Rg interactions do not affect Rg-Rg interactions [8]; no modifications have been made to the Rg-Rg potentials to account for perturbations arising from the presence of the M atom.

Rg-Rg AND M/Rg POTENTIALS

The Rg-Rg interaction potential in Eq. (3) can be approximated as a simple sum over the appropriate Σ symmetry Rg-Rg diatomic potentials

$$V_{\text{Rg-Rg}}(\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = \sum_{m=1, N} \sum_{m < k} V_{\text{Rg-Rg}}(|\mathbf{R}_m - \mathbf{R}_k|) \\ = U_{\text{Rg-Rg}} \quad (4)$$

following the notation of Boatz and Fajardo [7]. It should be noted that $U_{\text{Rg-Rg}}$ has the same value for both the 2S and 2P terms at this level of approximation.

The M-Rg potential, which in general exhibits a complicated dependence on the Rg positions (\mathbf{R}_k), is expressed in terms of a Legendre polynomial expansion as

$$V_{\text{M-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) \\ = \sum_{k=1, N} \sum_{L=0, \infty} V_L(\mathbf{r}, \mathbf{R}_k) P_L(\theta, \phi; \theta_k, \phi_k), \quad (5)$$

where $\mathbf{r} = (r, \theta, \phi)$ is the optically active electron position vector, and $\mathbf{R}_k = (R_k, \theta_k, \phi_k)$ is the position vector for Rg nucleus k . $V_L(\mathbf{r}, \mathbf{R}_k)$ is the radial potential function of order L , and $P_L(\theta, \phi; \theta_k, \phi_k)$ is the Legendre polynomial of order L

$$P_L(\theta, \phi; \theta_k, \phi_k) = 4\pi(2L+1)^{-1} \\ \times \sum_{M=-L, L} Y_{LM}(\theta, \phi) Y_{LM}^*(\theta_k, \phi_k) \quad (6)$$

and the Y_{LM} are spherical harmonic functions expressed in the standard Condon and Shortley phase convention [22]. Equation (6) is an expression of one of the possible forms of the spherical harmonic addition theorem [21].

SPIN-ORBIT COUPLING OPERATOR

The spin-orbit coupling operator for the optically active electron may be written, using the standard defining equation for the spin-orbit interaction [23], in terms of the potential gradient of the free M and M-Rg interaction potentials

$$H_{\text{SO}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = H_{\text{SO,M}}(\mathbf{r}) \\ + H_{\text{SO,M-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = \{(2m_e^2 c^2 r)^{-1} \nabla[V_M(\mathbf{r}) \\ + V_{\text{M-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N)] \times \mathbf{p}\} \cdot \mathbf{S} \\ = [\xi_M(r) + \xi_{\text{M-Rg}}(r)] \mathbf{L} \cdot \mathbf{S} + [(2m_e^2 c^2 r)^{-1} \\ \times \nabla \left\{ \sum_{k=1, N} \sum_{L=1, \infty} V_L(\mathbf{r}, \mathbf{R}_k) P_L(\theta, \phi; \theta_k, \phi_k) \right\} \times \mathbf{p}] \cdot \mathbf{S} \quad (7)$$

where \mathbf{p} and \mathbf{S} are, respectively, the linear momentum and spin operators for the optically active electron, $\mathbf{L} = \mathbf{r} \times \mathbf{p}$ is the angular momentum operator for the optically active electron, and

$$\xi_M(r) = (2m_e^2 c^2 r)^{-1} d[V_M(\mathbf{r})]/dr \quad (8)$$

is the standard spin-orbit coupling operator for the free M atom. The M-Rg interaction potential of Eq. (5) can be broken up into spherical and nonspherical components

$$V_{\text{M-Rg}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = \sum_{k=1, N} V_0(\mathbf{r}, \mathbf{R}_k) \\ + \sum_{k=1, N} \sum_{L=1, \infty} V_L(\mathbf{r}, \mathbf{R}_k) P_L(\theta, \phi; \theta_k, \phi_k). \quad (9)$$

In the form of the theory currently in use for MC-MCD simulations, it is assumed that the gradient expression in Eq. (7) is small and can be neglected; i.e., no $L > 0$ terms are assumed to contribute to the spin-orbit coupling interaction. The total spin-orbit coupling operator of the M/Rg system in the spherical approximation is thus given by

$$H_{\text{SO}}(\mathbf{r}) = H_{\text{SO,M}}(\mathbf{r}) + H_{\text{SO,M-Rg}}(\mathbf{r}) \\ = [\xi_M(r) + \xi_{\text{M-Rg}}(r)] \mathbf{L} \cdot \mathbf{S} = \xi(r, \mathbf{L} \cdot \mathbf{S}), \quad (10)$$

where the dependence on the Rg position vectors $\{\mathbf{R}_k\}$, while present, is not explicitly shown in the interest of preserving a compact notation in the final form of the equations used.

No spin-orbit coupling is possible when the optically active electron is in the 2S ground state manifold, where $l = 0$. In the 2P excited state manifold, where $l = 1$, the spin-orbit coupling constant is expressed formally as the sum of the radial expectation values of the spherically symmetric M and M-Rg spin-orbit coupling operators

$$\zeta_{nl} = \langle n l | \xi(r) | n l \rangle = \langle n l | \xi_M(r) | n l \rangle \\ + \langle n l | \xi_{\text{M-Rg}}(r) | n l \rangle = \zeta_M + \zeta_{\text{M-Rg}}. \quad (11)$$

If the radial eigenvector $|n l\rangle$ is the free atom P state atomic radial eigenvector for M, or if $|n l\rangle$ closely approximates this eigenvector, then ζ_M is the free alkali atom spin-orbit coupling constant and $\zeta_{\text{M-Rg}}$ is the spherical perturbation to the spin-orbit coupling constant induced in M by the Rg matrix environment. Equation (11) provides an insightful

context for discussing the possibilities of (1) spin-orbit coupling constant sign reversal or (2) reduction of the magnitude of a spin-orbit coupling constant (partial quenching), in a 2P M-Rg system. For any given alkali metal M, the free atom 2P spin-orbit coupling constant is positive; i.e., $\zeta_M > 0$. Thus, for a 2P spin-orbit coupling constant ζ_{nl} to be negative in a M-Rg system, the M-Rg spin orbit contribution must be negative and larger in magnitude than the free M atom spin-orbit coupling constant; i.e., $\zeta_{M-RG} < 0$ and $|\zeta_{M-RG}| > \zeta_M$. The conditions for the partial quenching case, where ζ_{nl} is still positive but smaller than ζ_M , are: $\zeta_{M-RG} < 0$ and $|\zeta_{M-RG}| < \zeta_M$. In the actual MC-MCD simulations, ζ_{nl} is introduced as an adjustable parameter. The two cases may be explored in the MC-MCD simulations by making appropriate choices for ζ_{nl} with respect to the known positive ζ_M of the alkali metal M atom of interest.

ZEEMAN PERTURBATION

The Zeeman perturbation operator is given by

$$H_{\text{Zeeman}}(\mathbf{r}; \mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) = \mu_B \mathbf{B}_{\text{local}}(\mathbf{L} + 2\mathbf{S}), \quad (12)$$

where μ_B is the Bohr magneton, \mathbf{L} and \mathbf{S} are the orbital and spin angular momentum operators for the optically active electron, and $\mathbf{B}_{\text{local}}$ is the magnetic field locally experienced by the optically active electron as a result of the application of an external magnetic field \mathbf{B} to the M/Rg system. Both the magnitude and the direction of the local magnetic field vector are influenced by the positions of the Rg atoms and by the nature of the electronic state of the optically active electron (i.e., 2S or 2P). The relationship between the external magnetic field and the local magnetic field thus may be expressed as

$$\mathbf{B}_{\text{local}} = \mathbf{B}_{\text{local}}(\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N) \\ = [1 - \sigma_{nl}(\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N)]\mathbf{B}, \quad (13)$$

where $\sigma_{nl}(\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N)$ is a 3×3 shielding tensor whose individual components depend on both the nature of the electronic state being shielded and the positions of the Rg atoms in the matrix. In actual MC-MCD simulations, it is assumed that the local magnetic field is aligned along the laboratory z-axis

$$\mathbf{B}_{\text{local}} = B_{\text{local}}\mathbf{e}_z, \quad (14)$$

which gives

$$H_{\text{Zeeman}} = \mu_B B(1 - \sigma_{nl})(L_z + 2S_z) \quad (15)$$

Different values of B_{local} can be chosen, if desired, to model the effects of slightly different 2S and 2P shielding constants σ_{n0} and σ_{n1} . Changes in the direction of B_{local} away from the z-axis arising from the tensor character of σ_{nl} can be modeled by preserving the laboratory z-axis orientation of the Zeeman perturbation, which is very desirable from the standpoint of matrix element evaluation, and rotating the Rg position vector set through an arbitrary unitary rotation transformation $U\{\mathbf{R}_k\} = \{\mathbf{R}'_k\}$. In any event, it is expected that differences between \mathbf{B} and $\mathbf{B}_{\text{local}}$ will be comparatively small for closed shell Rg atoms in their ground 1S electronic states. Certainly within the chosen MC-MCD simulation restrictions mentioned earlier of neglecting polarization effects and the higher excited states of the Rg atoms, \mathbf{B} will not differ much from $\mathbf{B}_{\text{local}}$.

DIAGONALIZATION OF THE FULL HAMILTONIAN MATRIX IN THE 2S AND 2P MANIFOLDS

An implementation of degenerate first-order perturbation theory is required to arrive at approximate 2S and 2P eigenvalues and eigenvectors of the full M/Rg Hamiltonian given in Eq. (1). These eigenvalues and eigenvectors contain the effects of the various terms in the interaction Hamiltonian Eq. (3). First-order perturbation theory gives eigenvalues that are correct through first order. The associated eigenvectors are those particular zeroth-order eigenvectors, expressed as orthonormal linear combinations of the zeroth-order eigenvector set $\{|n l m_l s m_s\rangle\}$,

$$\Psi_i = |n l i\rangle = \sum_{m_l = -l, l} \sum_{m_s = -1/2, 1/2} C_i^{nlm_lsm_s} |n l m_l s m_s\rangle, \quad (16)$$

$$\langle n l i' | n l i \rangle = \delta(i', i), \quad (17)$$

that diagonalize the 2×2 (2S manifold) or 6×6 (2P manifold) matrix formed by sandwiching the full Hamiltonian, Eq. (1), within the same uncoupled zeroth-order eigenvector set $\{|n l m_l s m_s\rangle\}$; i.e.,

$$\begin{aligned} \langle n'l'm'_s'm'_s|H|n\ l\ m_l\ s\ m_s\rangle \\ = \langle n'l'm'_s'm'_s|H_M|n\ l\ m_l\ s\ m_s\rangle \\ + \langle n'l'm'_s'm'_s|H_{\text{int}}|n\ l\ m_l\ s\ m_s\rangle. \end{aligned} \quad (18)$$

Both the first-order eigenvalues and the associated zeroth-order eigenvectors are needed for MC-MCD simulations. Since the first matrix element on the right side of Eq. (18) is the zeroth-order matrix element of the free M Hamiltonian, it is automatically diagonal in either the original zeroth-order basis $\{|nlm_lsm_s\rangle\}$

$$\begin{aligned} \langle n'l'm'_s'm'_s|H_M|n\ l\ m_l\ s\ m_s\rangle \\ = \varepsilon_{nl}\delta(n', n)\delta(l', l)\delta(m'_l, m_l)\delta(s', s)\delta(m'_s, m_s) \end{aligned} \quad (19)$$

or the new zeroth-order basis $\{|n\ li\rangle\}$

$$\langle n\ li'|H_M|n\ li\rangle = \varepsilon_{nl}\delta(n', n)\delta(l', l)\delta(i', i). \quad (20)$$

Within the 2S manifold, the interaction matrix on the right side of Eq. (18) also is diagonal; thus

$$\Psi_i(^2S) = |n\ 0\ i\rangle = \sum_{m_s=-1/2, 1/2} C_i^{n00m_s} |n\ 0\ 0\ \frac{1}{2}\ m_s\rangle, \quad (21)$$

which reduces to

$$\Psi_1(^2S) = |n\ 0\ 0\ \frac{1}{2}\ \frac{1}{2}\rangle, \quad (22a)$$

$$\Psi_2(^2S) = |n\ 0\ 0\ \frac{1}{2}\ -\frac{1}{2}\rangle. \quad (22b)$$

The i index is mapped onto the m_s index one-to-one in this specific case. The final expressions for the 2S eigenvalues, correct through first order, are obtained by collecting together the matrix elements previously derived for the Rg-Rg, M-Rg, and Zeeman perturbations:

$$\begin{aligned} E_i(^2S) &= \varepsilon_{n0}(^2S) + E_{n0i}(^2S) = \varepsilon_{n0}(^2S) + U_{\text{Rg-Rg}}(^2S) \\ &+ U_{\text{M-Rg}}(^2S) + U_{\text{Zeeman}}(^2S) \\ &= \varepsilon_{n0}(^2S) + \sum_{m=1,N} \sum_{m<k} V_{\text{Rg-Rg},X\Sigma}(|\mathbf{R}_m - \mathbf{R}_k|) \\ &+ \sum_{m=1,N} \sum_{m<k} V_{\text{M-Rg},X\Sigma}(R_k) + \mu_B B_{\text{local}}(2m_s). \end{aligned} \quad (23)$$

In Eq. (23), the eigenvalue convention of Boatz and Fajardo [7] is adopted: E_i is an eigenvalue of a specific 2S state Ψ_i correct through first order, ε_{n0} is the zeroth order 2S eigenvalue for the optically active electron of the free M atom, and E_{n0i} is the first-order eigenvalue correction term arising from the interaction Hamiltonian.

To obtain 2P final state eigenvalues correct through first order

$$E_f(^2P) = \varepsilon_{n1}(^2P) + E_{n1f}(^2P), \quad (24)$$

the first-order eigenvalue set ($E_{n1f}(^2P); f = 1, 6$) must be calculated by diagonalizing the 6×6 2P interaction matrix in Eq. (18). The individual matrix elements of this interaction matrix, shown in Table I, are obtained by collecting together the matrix elements previously derived for the 2P Rg-Rg, M-Rg, spin-orbit, and Zeeman perturbations. It should be noted that the $U_{\text{Rg-Rg}}$ and the $\langle V_0 \rangle_{\text{M-Rg}}$ perturbations are diagonal perturbations in this 6×6 matrix and can be separated from the 6×6 matrix eigenvalue problem as illustrated by Boatz and Fajardo [7] in their Eqs. (6) and (8). The associated final state 2P eigenvectors that diagonalize this matrix, are

$$\begin{aligned} \Psi_f(^2P) &= |n\ 1\ f\rangle \\ &= \sum_{m_1=-1, 1} \sum_{m_s=-1/2, 1/2} C_f^{n1m_1(1/2)m_s} |n\ 1\ m_1\ \frac{1}{2}\ m_s\rangle, \end{aligned} \quad (25)$$

where $f = 1, 6$ and

$$\langle n\ 1\ f'|n\ 1\ f\rangle = \delta(f', f). \quad (26)$$

$^2S \rightarrow ^2P$ TRANSITION ENERGIES AND TRANSITION MOMENT INTEGRALS FOR M/Rg MCD SPECTROSCOPY

The combined effects of the Rg-Rg, M-Rg, spin-orbit, and Zeeman perturbations on the optically active electron of a M/Rg system act to split the ground 2S manifold into two energetically distinct states $\{\Psi_i; i = 1, 2\}$ and the excited 2P manifold into six energetically distinct states $\{\Psi_f; f = 1, 6\}$. The specific characteristics of the 2S and 2P eigenvalues and their associated eigenvectors are dependent upon the particular Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}$ within which the problem is solved. Thus, for a given Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}$ and magnitude of the magnetic field, up to 12 energetically and optically distinct transitions may be identified

TABLE I
Matrix elements of the 6×6 2P interaction matrix in the uncoupled angular momentum representation $|nlm_s m_s\rangle$.

$ n11 \frac{1}{2} \frac{1}{2}\rangle$	$ n11 \frac{1}{2} \frac{1}{2}\rangle$	$ n10 \frac{1}{2} \frac{1}{2}\rangle$	$ n1 -1 \frac{1}{2} \frac{1}{2}\rangle$	$ n11 \frac{1}{2} -\frac{1}{2}\rangle$	$ n10 \frac{1}{2} -\frac{1}{2}\rangle$	$ n1 -1 \frac{1}{2} -\frac{1}{2}\rangle$
$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [-3 \cos^2 \theta_k - 1] + \frac{1}{2} \zeta_{nl} + 2\mu_B B + U_{Rg-Rg}$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3 \sin^2 \theta_k e^{-i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3\sqrt{2} \sin \theta_k \cos \theta_k e^{-i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3 \sin^2 \theta_k e^{-2i\phi_k}]$			
$\langle n10 \frac{1}{2} \frac{1}{2} $	$\frac{1}{10} \langle V_2 \rangle_k \times [-3\sqrt{2} \sin \theta_k \cos \theta_k e^{i\phi_k}]$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times 2(3 \cos^2 \theta_k - 1) + \mu_B B + U_{Rg-Rg}$	$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{-i\phi_k}]$	$\frac{1}{\sqrt{2}} \zeta_{nl}$		
$\langle n1 -1 \frac{1}{2} \frac{1}{2} $	$\frac{1}{10} \langle V_2 \rangle_k \times [-3 \sin^2 \theta_k e^{2i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{i\phi_k}]$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [-3 \cos^2 \theta_k - 1] - \frac{1}{2} \zeta_{nl} + U_{Rg-Rg}$	$\frac{1}{\sqrt{2}} \zeta_{nl}$		
$\langle n11 \frac{1}{2} -\frac{1}{2} $		$\frac{1}{\sqrt{2}} \zeta_{nl}$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [-3 \cos^2 \theta_k - 1] - \frac{1}{2} \zeta_{nl} + U_{Rg-Rg}$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3\sqrt{2} \sin \theta_k \cos \theta_k e^{-i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3\sqrt{2} \sin \theta_k \cos \theta_k e^{-2i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3 \sin^2 \theta_k e^{-2i\phi_k}]$
$\langle n10 \frac{1}{2} -\frac{1}{2} $		$\frac{1}{\sqrt{2}} \zeta_{nl}$	$\frac{1}{10} \langle V_2 \rangle_k \times [-3\sqrt{2} \sin \theta_k \cos \theta_k e^{i\phi_k}]$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [2(3 \cos^2 \theta_k - 1) - \mu_B B + U_{Rg-Rg}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{-i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{-2i\phi_k}]$
$\langle n1 -1 \frac{1}{2} -\frac{1}{2} $			$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{-i\phi_k}]$	$\frac{1}{10} \langle V_2 \rangle_k \times [3\sqrt{2} \sin \theta_k \cos \theta_k e^{i\phi_k}]$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [-3 \cos^2 \theta_k - 1] + \mu_B B + U_{Rg-Rg}$	$\langle V_0 \rangle_k + \frac{1}{10} \langle V_2 \rangle_k \times [-3 \cos^2 \theta_k - 1] + \frac{1}{2} \zeta_{nl} + 2\mu_B B + U_{Rg-Rg}$

^a $\langle V_0 \rangle_k = \frac{1}{3} [V_{B\pm}(R_k) + 2V_{\lambda\pi\pi}(R_k)]$ and $\langle V_2 \rangle_k = \frac{5}{3} [V_{B\pm}(R_k) - V_{\lambda\pi\pi}(R_k)]$. The final matrix elements are constructed by summing over "k", which indexes the rare gas atoms.

within the manifold-to-manifold $^2S \rightarrow ^2P$ transition of the M/Rg system. The transition energies of these 12 $^2S \rightarrow ^2P$ transitions are given as

$$h\nu_{fi} = E_f(^2P) - E_i(^2S), \quad (27)$$

where $i = 1, 2$ and $f = 1, 6$. The Rg–Rg interaction energy terms cancel in Eq. (27), since $U_{\text{Rg-Rg}}(^2P) = U_{\text{Rg-Rg}}(^2S) = U_{\text{Rg-Rg}}$.

Electronic transitions involving the optically active electron of a M/Rg system are well described as electric dipole transitions. In MCD spectroscopy, each of the 12 transitions may be induced to a greater or lesser extent by either LCP or RCP light interacting with the M/Rg system in the electric dipole limit and tuned to the particular transition energy $h\nu_{fi}$. Thus, both LCP and RCP electric dipole transition moment integrals must be computed for each of the 12 possible combinations of initial and final states; i.e., 24 transition moment integrals, 12 for LCP light and 12 for RCP light, are needed to describe the MCD spectrum of a $^2S \rightarrow ^2P$ transition in M/Rg system in a particular Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}$; i.e.,

$$\langle \Psi_f(^2P) | O_{\text{LCP}} | \Psi_i(^2S) \rangle \quad (28a)$$

$$\langle \Psi_f(^2P) | O_{\text{RCP}} | \Psi_i(^2S) \rangle, \quad (28b)$$

where $i = 1, 2$ and $f = 1, 6$. O_{LCP} and O_{RCP} are the electric dipole transition operators for LCP and RCP light in Eqs. (28a) and (28b). The electric dipole moment operator \mathbf{m} of the M/Rg system is, to a very good initial approximation, determined by the position $\mathbf{r} = (x, y, z)$ and charge $-e$ of the optically active electron. The Rg atoms are electrically neutral, and their contributions to the dipole moment can be neglected if polarization effects are assumed to be small. The M^+ core is positioned at the origin of the chosen coordinate system and thus does not contribute to the expression for the dipole moment at this level of approximation, i.e.,

$$\begin{aligned} \mathbf{m} &= \sum_i q_i \mathbf{r}_i = -e\mathbf{r} + e\mathbf{0} = -e(x\mathbf{e}_x + y\mathbf{e}_y + z\mathbf{e}_z) \\ &+ e(0\mathbf{e}_x + 0\mathbf{e}_y + 0\mathbf{e}_z) = -e(x\mathbf{e}_x + y\mathbf{e}_y + z\mathbf{e}_z). \end{aligned} \quad (29)$$

However, the optically active electron at position \mathbf{r} will induce a slight polarization in the M^+ core and the Rg matrix. This polarization can be represented as a small negative electrostatic image

charge $-q'$ at $-\mathbf{r}$. Thus, polarization effectively reduces the charge of the optically active electron to

$$-e' = -e + q', \quad (30)$$

where $|-e'| < |-e|$, and the dipole moment operator of the M/Rg system becomes

$$\mathbf{m} = -e'\mathbf{r} = -e'(x\mathbf{e}_x + y\mathbf{e}_y + z\mathbf{e}_z). \quad (31)$$

While it can be argued that the image charge $-q'$ (and hence $-e'$) is not, strictly speaking, a constant, but will exhibit both a radial dependence (i.e., different q' for small r vs. large r) and angular dependence arising from the specific positions of the Rg atoms ($\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N$) in the M/Rg system, these effects are neglected at this level of the theory. Within the Piepho and Schatz unit vector conventions for LCP and RCP light [9],

$$\pi_{\text{LCP}} = 2^{-1/2}(\mathbf{e}_x - i\mathbf{e}_y) \quad (32a)$$

$$\pi_{\text{RCP}} = 2^{-1/2}(\mathbf{e}_x + i\mathbf{e}_y), \quad (32b)$$

the LCP and RCP electric dipole moment operators for the M/Rg system become

$$\begin{aligned} O_{\text{LCP}} &= \mathbf{m} \cdot \pi_{\text{LCP}}^* = -e'2^{-1/2}(x + iy) \\ &= e'(4\pi/3)^{1/2}Y_{11}(\theta, \phi)r, \end{aligned} \quad (33a)$$

$$\begin{aligned} O_{\text{RCP}} &= \mathbf{m} \cdot \pi_{\text{RCP}}^* = -e'2^{-1/2}(x - iy) \\ &= -e'(4\pi/3)^{1/2}Y_{1-1}(\theta, \phi)r. \end{aligned} \quad (33b)$$

Since each of the $\Psi_i(^2S)$ and $\Psi_f(^2P)$ states is expressed in terms of the uncoupled zeroth-order angular momentum eigenvector set $\{|nlm_l m_s\rangle\}$ [see Eqs. (16) and (25)], the specific form of the transition moment integrals given in Eqs. (33a) and (33b) may be determined by evaluating integrals

$$\begin{aligned} &\langle n l' m'_l s' m'_s | O_{\text{LCP}} | n l m_l m_s \rangle \\ &= \langle n l' m'_l \tfrac{1}{2} m'_s | O_{\text{LCP}} | n 0 0 \tfrac{1}{2} m_s \rangle = f\delta(m'_l, 1)\delta(m'_s, m_s), \end{aligned} \quad (34a)$$

$$\begin{aligned} &\langle n l' m'_l s' m'_s | O_{\text{RCP}} | n l m_l m_s \rangle \\ &= \langle n l' m'_l \tfrac{1}{2} m'_s | O_{\text{RCP}} | n 0 0 \tfrac{1}{2} m_s \rangle \\ &= -f\delta(m'_l, -1)\delta(m'_s, m_s), \end{aligned} \quad (34b)$$

where

$$f_{\text{LCP}} = f = (e'[4\pi/3]^{1/2})\langle n1|r|n0\rangle\langle Y_{11}|Y_{00}\rangle \\ = (e'[3]^{-1/2})\langle n1|r|n0\rangle \quad (35a)$$

$$f_{\text{RCP}} = -f = (-e'[4\pi/3]^{1/2})\langle n1|r|n0\rangle\langle Y_{1-1}|Y_{00}\rangle \\ = (e'[3]^{-1/2})\langle n1|r|n0\rangle \quad (35b)$$

Equations (34a) and (35a) express the fact that an LCP electric dipole transition acts on the orbital angular momentum component of the uncoupled wave function, increasing both l and m_l by one unit while leaving s and m_s unchanged. The corresponding RCP electric dipole transition described in Eqs. (34b) and (35b) also acts on the orbital angular momentum in the uncoupled wave function, increasing l by one unit and decreasing m_l by one unit. In Eqs. (35a) and (35b), f is the magnitude of the evaluated transition moment integral. Note that the value of f is the same for all allowed transitions. By combining the transition moment integral expressions in Eqs. (28), (34), and (35) with the appropriate expressions for $\Psi_i(^2S)$ and $\Psi_f(^2P)$ in the uncoupled basis as given in Eqs. (16), (17), and (25), final expressions for the contribution to the MCD spectrum by the transition $i \rightarrow f$ at energy $h\nu_{fi}$ can be written:

$$\Delta A'(v_{fi}) = A_{\text{LCP}}(v_{fi}) - A_{\text{RCP}}(v_{fi}) \\ = \alpha\{|\langle \Psi_f(^2P)|O_{\text{LCP}}|\Psi_i(^2S)\rangle|^2 - |\langle \Psi_f(^2P)|O_{\text{RCP}}|\Psi_i(^2S)\rangle|^2\} \\ = \alpha f^2\{|C_f^{n1,m_l=1,m_s}|^2 - |C_f^{n1,m_l=-1,m_s}|^2\} \\ \times [1 + (e^{-2\mu_B B/kT} - 1)\delta(m_s, 1/2)], \quad (36)$$

where α is a numerical constant that relates the squared transition moment integrals to the absorbances and the initial state index i is associated with $m_s = -1/2$ or $1/2$ in the 2S manifold. The MCD spectral amplitude at energy $h\nu_{fi}$ for the Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}$ is seen to be related to differences of squares of selected 2P eigenvector coefficients computed for that configuration. The term in square brackets on the right side of Eq. (36) is a Boltzmann factor for the 2S manifold. The Zeeman perturbation splits the energies of the $m_s = -1/2$ and $1/2$ states of the 2S manifold by a factor of $2\mu_B B$. The $m_s = -1/2$ and $1/2$ states will, therefore, have different thermal populations. This must be accounted for in computing the various contributions to the MCD spectrum.

Monte Carlo Simulation of the MCD Spectrum

The full MCD spectrum for a $^2S \rightarrow ^2P$ transition in a M/Rg system in a specific Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}$ is, therefore, a "stick MCD spectrum" consisting of 12 lines, appropriately placed on the x (energy) axis at transition energies $h\nu_{fi}$, whose magnitudes and directions (+ or -) on the y ($\Delta A'$) axis are determined by computing Eq. (36) for all possible choices of i and f . To simulate a real MCD spectrum, many stick MCD spectra, each arising from a different Rg configuration, are averaged together using a simple binning (histogram) technique based on the original Metropolis et al. [24] algorithm exactly as implemented by Boatz and Fajardo [7] for the case of electronic absorption spectroscopy. A MC-MCD simulation begins by choosing an initial Rg configuration $\{\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N\}_{\text{initial}}$, typically, representing an idealized trapping site (e.g., a single substitutional site with O_h symmetry). The MC energy optimization scheme operates in the 2S ground state manifold. At this stage, the very small Zeeman perturbation of the 2S manifold is neglected. MCD stick spectra arising from more favorable configurations are weighted more heavily in the averaging process than are those arising from the less favorable configurations. In actual practice, the LCP and RCP contributions to the MCD spectrum appearing in Eq. (36) are accumulated and stored separately. This allows the electronic absorption spectrum to be recovered from the MC-MCD simulation as [9]

$$A' = A = \frac{1}{2}[A'_{\text{LCP}} + A'_{\text{RCP}}]. \quad (37)$$

Equation (37) expresses the fact that the electronic absorption spectra of M/Rg systems are essentially unaffected by the application of an external magnetic field.

ORIENTATIONAL AVERAGING IN MC-MCD SIMULATIONS

In an MCD experiment, the propagation direction of the light, which is always parallel to the \mathbf{B} vector, rigorously defines the laboratory frame z -axis. The laboratory frame x - and y -axes are set parallel to the sapphire deposition window and perpendicular to \mathbf{B} with the origin at the M nucleus. All Rg positions, eigenvectors, and selection rules

are described with respect to this laboratory frame. To model the effects of arbitrary Rg lattice orientation relative to the laboratory frame, the 6×6 2P matrix listed in Table I is separated into M/Rg and spin-orbit + Zeeman matrices:

$$[M/Rg + SO + Zeeman] = [M/Rg] + [SO + Zeeman]. \quad (38)$$

Arbitrary orientations of the Rg matrix relative to the laboratory frame can be modeled by rewriting Eq. (38) as

$$[M/Rg + SO + Zeeman] = [U][M/Rg][U]^{-1} + [SO + Zeeman], \quad (39)$$

where U is an arbitrary randomly generated unitary rotation matrix (e.g., a Eulerian rotation matrix). The MC-MCD simulation process is carried out as before, using Eq. (39) rather than Eq. (38).

LATTICE TEMPERATURE AND MAGNETIC TEMPERATURE IN MC-MCD SIMULATIONS

Temperature effects appear in two distinct places in MC-MCD simulations. A lattice temperature must be specified for the MC simulation process. It may be argued that a proper representation of quantum mechanical zero point vibrational motions of the M/Rg lattice at temperature T requires the classical MC simulation to be carried out at a classical lattice temperature $T' > T$ [8]. However, a temperature for the Boltzmann factor of the Zeeman-split 2S term also must be included. This magnetic temperature, T_{mag} , should always be the actual system temperature T ($T_{\text{mag}} = T$). The MC-MCD simulation code has the option of specifying separate values for the lattice temperature T' and the magnetic temperature T_{mag} . Lawrence and Apakarian [8] discuss the relationship between T' and T in classical MC simulations of doped Rg systems. In practice T' is calibrated to T by comparing the Rg MC equilibrium lattice spacing (for first and second nearest neighbors) with the experimentally measured values [25]. For Ar at 10 K, the first nearest-neighbor analysis predicts a lattice temperature of 31 K, while the second nearest-neighbor analysis predicts a slightly lower lattice temperature of 27 K. Consequently a simulation lattice temperature in the vicinity of 30 K represents the best modeling of the quantum zero point vibrations of the Ar lattice. The Na/Ar system at a magnetic temperature,




FIGURE 1. Comparison of the temperature effects in the (a) absorption, and (b) MCD spectra for the case of $SO = 17 \text{ cm}^{-1}$, $B = 1.0 \text{ T}$, magnetic temperature, $T_m = 10 \text{ K}$, at simulation temperatures of 10, 30, and 45 K.

$T_{\text{mag}} = 10 \text{ K}$, a Na 2P spin-orbit parameter of 17 cm^{-1} (the value for unperturbed atomic Na) is illustrated in Figure 1(a, b), in which are presented the simulated absorption and MCD spectra, respectively, for three different choices of lattice temperature, $T' = 10, 30$, and 45 K .




FIGURE 2. Absorption spectra with (a) normal and (b) reversed B-fields. 1-atom vacancy, $SO = 17 \text{ cm}^{-1}$, and $T_{\text{mag}} = 6, 8, 10 \text{ K}$. The magnitudes of the B-fields are 1.0 T.

While the characteristic three-peaked “triplet” absorption feature and the corresponding down–down–up MCD triplet feature are preserved for all three choices of T' , the spectral breadth increases as T' increases, improving agreement between simulated and experimental spectra. The difference in peak height is due to the fact that the total area under the absorption curve is normalized to 1.0; broader peaks will be shorter.

OTHER IMPORTANT PROPERTIES OF MC-MCD SIMULATIONS

For the single substitutional site in the Na/Ar system, absorption and MC-MCD simulations correctly reproduce other observed properties of ex-

perimental absorption and MCD spectra M/Rg systems. For example, the observed insensitivity of the experimental $^2S \rightarrow ^2P$ M/Rg triplet absorption feature to the magnetic field strength and T_{mag} is faithfully reproduced in the MC-MCD simulation, as shown in Figure 2. Reversal of the magnetic field direction ($\mathbf{B} = B\mathbf{e}_z$ changed to $\mathbf{B} = -B\mathbf{e}_z$) in an MCD experiment has the effect of inverting the MCD spectrum. If the normal MCD spectrum follows a down–down–up pattern, the MCD spectrum with the reversed magnetic field will be the mirror image of the original spectrum with an up–up–down pattern. Figure 3 shows that the MC-MCD simulations of Na/Ar correctly give an inverted spectrum when \mathbf{B} is replaced with $-\mathbf{B}$. These successes suggest the




FIGURE 3. MCD spectra with (a) normal magnetic field alignment, and (b) reversed magnetic field alignment. 1-atom vacancy, $SO = 17 \text{ cm}^{-1}$, and $T_{\text{mag}} = 10 \text{ K}$. The magnitudes of the B-fields are 1.0 T.

MC-MCD simulation embodies much of the essential physics of the Na/Ar system and its interaction with external magnetic fields, the perturbing effects of the Rg matrix and electromagnetic radiation. Overall, the MC-MCD simulated Na/Ar absorption and MCD spectra show good qualitative agreement with experimental Na/Ar spectra.

ACKNOWLEDGMENTS

It is a great pleasure to acknowledge Mario E. Fajardo for his helpful discussions on theory and for his insight in making connections between theory and experiment.

References

1. Fajardo, M. E.; Carrick, P. G.; Kenney, J. W. III. *J Chem Phys* 1991, 94, 5812.
2. Fajardo, M. E. *J Chem Phys* 1993, 98, 110.
3. Tam, S.; Fajardo, M. *J Chem Phys* 1993, 99, 854.
4. Corbin, R.; Fajardo, M. *J Chem Phys* 1994, 101, 2678.
5. Thompson, T. L.; Cordonnier, M. E.; Fajardo, M. E.; Tam, S. *Chem Phys* 1994, 189, 361.
6. Carrick, P. G. Specific impulse calculations of high energy density solid cryogenic rocket propellants. I. Atoms in solid h2. Technical Report PL-TR-93-3014; Phillips Laboratory: Edwards AFB, CA, 1993.
7. Boatz, J.; Fajardo, M. *J Chem Phys* 1994, 101, 3472.
8. Lawrence, W. G.; Apkarian, V. A. *J Chem Phys* 1994, 101, 1820.
9. Piepho, S. B.; Schatz, P. N. *Group Theory in Spectroscopy With Applications to Magnetic Circular Dichroism*; Wiley: New York, 1983.
10. Mort, J.; Luty, F. *Phys Rev Lett* 1964, 12, 45.
11. Henry, C. H.; Snatterly, S. E.; Schlickter, C. P. *Phys Rev A: At Mol Opt Phys* 1965, 137, A583.
12. Brown, F. C.; Mort, J.; Luty, F. *Phys Rev A: At Mol Opt Phys* 1965, 137, 566.
13. Merle d'Aubinge, Y.; Roussel, A. *Phys Rev B: Condensed Matter and Material Physics* 1971, 3, 1421.
14. Stephens, P. J.; Osborne, G. A. *J Chem Phys* 1972, 56, 609.
15. Hormes, J.; Schiller, J. *Chem Phys* 1983, 74, 433.
16. Lund, P. A.; Smith, D.; Jacobs, S. M.; Schatz, P. N. *J Phys Chem* 1984, 88, 31.
17. Rose, J.; Smith, D.; Williamson, B. E.; Schatz, P. N.; O'Brien, M. C. M. *J Phys Chem* 1986, 90, 2608.
18. Samet, C.; Rose, J. L.; Williamson, B. E.; Schatz, P. N. *Chem Phys Lett* 1987, 142, 557.
19. O'Brien, M. C. M. *J Phys C: Solid State Phys* 1985, 18, 4963.
20. O'Brien, M. C. M. *J Chem Phys* 1985, 82, 3870.
21. Weissbluth, M. *Atoms and Molecules*; Academic Press: San Diego, CA, 1978.
22. Condon, E. U.; Shortley, G. H. *The Theory of Atomic Spectra*; Cambridge: London, 1963.
23. Griffith, J. S. *Theory of Transition Metal Ions*; Cambridge: London, 1961.
24. Metropolis, N.; Rosenbluth, A. W.; Rosenbluth, M. N.; Teller, A. H.; Teller, E. *J Chem Phys* 1953, 21, 1087.
25. Terrill-Stolper, H. A. *Theory of the Magnetic Circular Dichroism (MCD) and Electronic Absorption Spectra of Alkali Metals Matrix Isolated in Cryogenic Rare Gases. I*; Master's thesis; New Mexico State University, Las Cruces, NM, 1996.